

THE DISCOVERY OF Y DWARFS USING DATA FROM THE *WIDE-FIELD INFRARED SURVEY EXPLORER* (WISE)

MICHAEL C. CUSHING^{1,10}, J. DAVY KIRKPATRICK², CHRISTOPHER R. GELINO², ROGER L. GRIFFITH², MICHAEL F. SKRUTSKIE³,
 A. MAINZER¹, KENNETH A. MARSH², CHARLES A. BEICHMAN², ADAM J. BURGASSER^{4,5,11}, LISA A. PRATO⁶,
 ROBERT A. SIMCOE⁵, MARK S. MARLEY⁷, D. SAUMON⁸, RICHARD S. FREEDMAN⁷,
 PETER R. EISENHARDT¹, AND EDWARD L. WRIGHT⁹

¹ Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, MS 321-520, Pasadena, CA 91109, USA; michael.cushing@gmail.com

² Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125, USA

³ Department of Astronomy, University of Virginia, Charlottesville, VA 22904, USA

⁴ Center for Astrophysics and Space Science, University of California San Diego, La Jolla, CA 92093, USA

⁵ Massachusetts Institute of Technology, 77 Massachusetts Avenue, Building 37, Cambridge, MA 02139, USA

⁶ Lowell Observatory, 1400 West Mars Hill Road, Flagstaff, AZ 86001, USA

⁷ NASA Ames Research Center, MS 254-3, Moffett Field, CA 94035, USA

⁸ Los Alamos National Laboratory, MS F663, Los Alamos, NM 87545, USA

⁹ Department of Physics and Astronomy, UCLA, Los Angeles, CA 90095, USA

Received 2011 June 10; accepted 2011 August 2; published 2011 November 22

ABSTRACT

We present the discovery of seven ultracool brown dwarfs identified with the *Wide-field Infrared Survey Explorer* (WISE). Near-infrared spectroscopy reveals deep absorption bands of H₂O and CH₄ that indicate all seven of the brown dwarfs have spectral types later than UGPS J072227.51–054031.2, the latest-type T dwarf currently known. The spectrum of WISEP J182831.08+265037.8 is distinct in that the heights of the *J*- and *H*-band peaks are approximately equal in units of f_{λ} , so we identify it as the archetypal member of the Y spectral class. The spectra of at least two of the other brown dwarfs exhibit absorption on the blue wing of the *H*-band peak that we tentatively ascribe to NH₃. These spectral morphological changes provide a clear transition between the T dwarfs and the Y dwarfs. In order to produce a smooth near-infrared spectral sequence across the T/Y dwarf transition, we have reclassified UGPS 0722–05 as the T9 spectral standard and tentatively assign WISEP J173835.52+273258.9 as the Y0 spectral standard. In total, six of the seven new brown dwarfs are classified as Y dwarfs: four are classified as Y0, one is classified as Y0 (pec?), and WISEP J1828+2650 is classified as >Y0. We have also compared the spectra to the model atmospheres of Marley and Saumon and infer that the brown dwarfs have effective temperatures ranging from 300 K to 500 K, making them the coldest spectroscopically confirmed brown dwarfs known to date.

Key words: brown dwarfs – infrared: stars – stars: individual (UGPS J072227.51–054031.2, WISEPC J014807.25–720258.8, WISEP J041022.71+150248.5, WISEPC J140518.40+553421.5, WISEP J154151.65–225025.2, WISEP J173835.52+273258.9, WISEP J182831.08+265037.8, WISEPC J205628.90+145953.3) – stars: low-mass

Online-only material: color figures

1. INTRODUCTION

Brown dwarfs, objects with too little mass to sustain the high core temperatures necessary for stable thermonuclear fusion of hydrogen, are the least massive, and possibly the most numerous products of star formation. Although first predicted to exist in the early 1960s (Kumar 1963; Hayashi & Nakano 1963), it was not until decades later that brown dwarfs were discovered in bulk by wide-area, red (700–1000 nm) and near-infrared (1–2.5 μ m) capable surveys such as the Two Micron All Sky Survey (2MASS; Skrutskie et al. 2006), the Sloan Digital Sky Survey (SDSS; York et al. 2000), and the Deep Near-Infrared Southern Sky Survey (DENIS; Epchtein et al. 1997). The emergent spectra of brown dwarfs are so distinct from that of late-type M dwarfs that the creation of two new spectral classes, L and T (Kirkpatrick et al. 1999; Burgasser et al. 2006), was required in order to properly classify them.¹²

The latest-type T dwarfs currently known were discovered in the UKIRT Infrared Deep Sky Survey (UKIDSS; Lawrence et al. 2007) and the Canada France Brown Dwarf Survey (CFBDS; Delorme et al. 2008b) and have estimated effective temperatures (T_{eff} s) of 500–700 K (e.g., Burningham et al. 2008; Delorme et al. 2008a; Leggett et al. 2009; Lucas et al. 2010; Liu et al. 2011).

Despite these spectacular successes, there exists a gap of nearly 400 K between the coolest spectroscopically confirmed brown dwarfs at $T_{\text{eff}} \sim 500$ K (Lucas et al. 2010) and Jupiter at $T_{\text{eff}} \sim 124$ K (Hanel et al. 1981). Although observations of star formation regions and young associations such as the Orion Nebula Cluster (Weights et al. 2009), Chameleon I (Luhman et al. 2005), and TW Hydrae (Chauvin et al. 2004) suggest that nature can form brown dwarfs that will eventually cool to these temperatures once they have dispersed from their natal clusters, they are simply too faint to have been detected by the aforementioned surveys. Recently however, two brown dwarfs with estimated effective temperatures of 300–400 K, WD 0806–661B (Luhman et al. 2011) and CFBDSIR J145829+101343B (CFBDSIR J1458+1013B; Liu et al. 2011), were discovered in targeted searches for companions to nearby stars. Although

¹⁰ Current address: Department of Physics and Astronomy, The University of Toledo, 2801 West Bancroft Street, Toledo, OH 43606, USA.

¹¹ Hellman Fellow.

¹² A compendium of known L and T dwarfs can be found at <http://DwarfArchives.org>.

efforts to obtain spectra of these two common proper motion sources have been hampered by extreme faintness in the case of WD 0806–661 ($J > 21.7$; Rodriguez et al. 2011), and proximity to its primary star in the case of CFBDSIR J1458+1013B ($<0''.11$), their mere existence suggests that a population of nearby cold brown dwarfs awaits discovery.

Foremost is the question of what these objects will look like spectroscopically and whether a new spectral class beyond T, dubbed “Y” (Kirkpatrick et al. 1999; Kirkpatrick 2000), will be required in order to properly classify them. Chemical equilibrium calculations and model atmospheres predict that as brown dwarfs cool below $T_{\text{eff}} \sim 600$ K, their atmospheres pass through a series of chemical transitions which in turn impact the appearance of their emergent spectra (Lodders 1999; Burrows & Sharp 1999; Lodders & Fegley 2002; Burrows et al. 2003). At $T_{\text{eff}} \sim 600$ K, the combination and overtone bands of NH_3 emerge in the near-infrared.¹³ At $T_{\text{eff}} \sim 500$ K, the prominent resonance absorption lines of Na I and K I in the red optical spectra of warmer brown dwarfs weaken as Na condenses out of the gas phase into Na_2S and then K condenses into KCl. Finally, H_2O and NH_3 will also condense out at $T_{\text{eff}} \sim 350$ K and ~ 200 K, respectively. Although each transition has been suggested as the trigger for the Y spectral class, focus has primarily been on detecting the NH_3 bands because they are predicted to emerge at the hottest effective temperatures. Although NH_3 absorption has been tentatively detected in the near-infrared spectrum of CFBDS J005910.90–011401.3 (hereafter CFBDS J0059–0114; Delorme et al. 2008a), this feature has not been confirmed in the spectrum of the cooler object UGPS J072227.51–054031.2 (UGPS 0722–05; Lucas et al. 2010).

Independent of their spectral morphology, the study of these ultracool brown dwarfs will provide important insights into both ultracool atmospheric physics and the low-mass end of the stellar mass function. Because brown dwarfs and exoplanets have similar atmospheric conditions, ultracool brown dwarfs are excellent exoplanet analogs that can be used as benchmarks for model atmospheres. The study of these ultracool brown dwarfs will therefore directly inform the interpretation and characterization of exoplanets detected with the next generation of high-contrast imagers like the Gemini Planet Imager (GPI; Macintosh et al. 2006), the Spectro-Polarimetric High-contrast Exoplanet Research (SPHERE) instrument for the Very Large Telescope (Beuzit et al. 2006), Project 1640 at Palomar Observatory (Hinkley et al. 2011), and the L - and M -band Infrared Camera (LMIRcam; Skrutskie et al. 2010) for the Large Binocular Telescope Interferometer (LBTI). Simulations by Burgasser (2004) and Allen et al. (2005) have also shown that the space density of cold brown dwarfs is very sensitive to both the underlying mass function and the low-mass limit of star formation. Identifying and characterizing a statistically robust sample of cold brown dwarfs will therefore provide two critical constraints on theories of low-mass star formation (Whitworth & Stamatellos 2006; Whitworth et al. 2007).

One of the primary science goals of the *Wide-field Infrared Survey Explorer* (WISE; Wright et al. 2010), a NASA mission that recently surveyed the entire sky at 3.4 (W1), 4.6 (W2), 12 (W3), and 22 (W4) μm , is to identify such cold brown dwarfs. The W1 and W2 bands were designed specifically to sample the

deep CH_4 absorption band centered at 3.3 μm and the region relatively free of opacity centered at $\sim 4.7 \mu\text{m}$ in the spectra of cold brown dwarfs (see Figure 2 of Mainzer et al. 2011). Since the peak of the Planck function at these low effective temperatures is in the mid-infrared, a large amount of flux emerges from the 4.7 μm opacity hole, making the W1–W2 colors extremely red ($W1-W2 > 2$; Mainzer et al. 2011; Kirkpatrick et al. 2011). Indeed such red colors are almost unique amongst astronomical sources making the identification of cool brown dwarfs with the W1–W2 color alone relatively easy (see Figure 12 of Wright et al. 2010).

We have been conducting a search for cold brown dwarfs since the start of the WISE survey in mid 2010 January. This search has already born fruit with the discovery of six late-type T dwarfs (Mainzer et al. 2011; Burgasser et al. 2011) two of which have spectral types later than T8. Kirkpatrick et al. (2011) present over 100 new brown dwarfs, the vast majority of which have spectral types later than T6. In this paper, we focus on seven of the ~ 100 brown dwarfs whose near-infrared spectra indicate that they are the latest-type spectroscopically confirmed brown dwarfs currently known. Indeed, we identify six of these brown dwarfs as the first members of the Y spectral class. In Section 2, we briefly discuss our selection criteria before presenting the ground- and space-based imaging and spectroscopic follow-up observations in Section 3. In Section 4, we present the properties of the first Y dwarfs, define the transition between the T sequence and the Y dwarfs, and derive estimates of the atmospheric parameters of the new brown dwarfs using model atmospheres.

2. CANDIDATE SELECTION

The seven new brown dwarfs were identified as part of a larger ongoing search for cold brown dwarfs using WISE. A detailed description of this survey and our search criteria is presented by Kirkpatrick et al. (2011). Briefly, candidates were selected from the Source Working database derived from the first-pass processing operational co-adds using color constraints derived from known T dwarfs and model atmospheres (in the case of brown dwarfs with $T_{\text{eff}} < 500$ K). Table 1 lists the WISE designations and photometry of the seven brown dwarfs, and Figure 1 shows $2' \times 2'$ DSS I , 2MASS J and H , WISE W1, W2, and W3, and W1W2W3 color composite images for each dwarf. Hereafter, we abbreviate the numerical portions of the WISE designations as hhmm±ddmm, where the suffix is the sexagesimal right ascension (hours and minutes) and declination (degrees and arcminutes) at J2000.0 equinox.

3. OBSERVATIONS

The follow-up ground- and space-based observations of the seven WISE brown dwarfs are discussed in the following sections. Although we present the near-infrared photometry of the brown dwarfs in this work for completeness, we defer the discussion of these data to Kirkpatrick et al. (2011) who present a more detailed discussion that places them in context with the larger population of brown dwarfs. In addition to the observations of the seven WISE brown dwarfs, we also obtained a near-infrared spectrum of UGPS 0722–05 for comparison purposes. A log of the near-infrared photometric observations as well as the resulting photometry is given in Table 2, and a log of the spectroscopic observations is given in Table 3.

¹³ Although the fundamental band of NH_3 has been detected in the spectra of warmer T dwarfs at 10.5 μm (e.g., Roellig et al. 2004; Cushing et al. 2006), the intrinsically weaker near-infrared bands require a higher NH_3 abundance, and thus lower effective temperature, to become a dominant opacity source.

Table 1
WISE Photometry

Object	W1 (mag)	W2 (mag)	W3 (mag)	W4 (mag)	W1–W2 (mag)
WISEPC J014807.25–720258.8	18.812 ± 0.529^a	14.584 ± 0.052	> 12.579	> 9.521	4.228 ± 0.532
WISEP J041022.71 + 150248.5	> 18.101	14.190 ± 0.059	12.472 ± 0.482^a	> 8.923	$> 3.911 \pm 0.059$
WISEPC J140518.40 + 553421.5	> 17.989	14.085 ± 0.041	12.312 ± 0.252	> 9.115	$> 3.904 \pm 0.041$
WISEP J154151.65–225025.2	> 17.018	13.982 ± 0.112	12.134 ± 0.443^a	> 9.064	$> 3.036 \pm 0.112$
WISEP J173835.52+273258.9	18.155 ± 0.362	14.535 ± 0.057	12.536 ± 0.350	> 9.182	3.620 ± 0.366
WISEP J182831.08+265037.8	> 18.452	14.276 ± 0.050	12.320 ± 0.291	9.147 ± 0.438^a	$> 4.176 \pm 0.050$
WISEPC J205628.90 + 145953.3	> 17.742	13.852 ± 0.043	11.791 ± 0.222	> 8.646	$> 3.890 \pm 0.043$

Notes. Objects designated as WISEP are from the Preliminary Release Source Catalog while objects designated as WISEPC are from the first-pass processing operations co-add Source Working Database. Magnitudes are in the Vega system and are based on profile fits (w1mpro, w2mpro, w3mpro, w4mpro). Upper limits are at the 95% confidence level (see http://wise2.ipac.caltech.edu/docs/release/prelim/expsup/sec4_3c.html#ul2).

^a $S/N \leq 3$.

Table 2
Near-infrared Photometry

Object	Filter	Instrument	Magnitude (mag)	Exposure Time (s)	Co-adds	Number of Images	Total Exp. (s)	Date (UT)
WISEPC J0148–7202	<i>J</i>	PANIC	18.96 ± 0.07	30	1	18	540	2010 Aug 1
	<i>H</i>	PANIC	19.22 ± 0.04	15	1	108	1620	2010 Aug 1
WISEP J0410 + 1502	<i>J</i>	WIRC	19.25 ± 0.05	60	1	15	900	2010 Aug 29
	<i>H</i>	WIRC	19.05 ± 0.09	30	4	15	1800	2010 Jul 26
WISEPC J1405 + 5534	<i>J</i>	WIRC	20.20 ± 0.13	30	2	15	1800	2010 Jul 26
	<i>H</i>	WIRC	21.45 ± 0.41	5400	Multiple
WISEP J1541–2250	<i>J</i>	NEWFIRM	21.16 ± 0.36	30	2	10	600	2011 Apr 17
	<i>H</i>	NEWFIRM	20.99 ± 0.52	5	12	10	600	2011 Apr 17
WISEP J1738+2732	<i>J</i>	WIRC	19.47 ± 0.08	60	1	15	900	2010 Jul 26
	<i>H</i>	WIRC	20.66 ± 0.38	30	2	15	900	2010 Jul 26
WISEP J1828+2650	<i>J</i>	NIRC2	23.57 ± 0.35	120	1	6	720	2010 Jul 1
	<i>H</i>	NIRC2	22.85 ± 0.24	120	1	9	1080	2010 Jul 1
WISEPC J2056 + 1459	<i>J</i>	WIRC	19.31 ± 0.12	60	1	15	900	2010 Aug 29
	<i>H</i>	WIRC	> 19.5	30	2	15	900	2010 Aug 29
	<i>J</i>	NIRC2	19.21 ± 0.07	120	1	9	1080	2010 Jul 1
	<i>H</i>	NIRC2	19.56 ± 0.18	120	1	6	720	2010 Jul 1

Table 3
Spectroscopy Log

Object	Instrument	UT Date	Mode	Slit Width (arcsec)	Int. Time (s)	A0 V Calibrator Star
WISEPC J0148–7202	FIRE/Magellan	2010 Sep 18	Longslit	0.6	960	HD 1881
WISEP J0410 + 1502	FIRE/Magellan	2010 Nov 18	Longslit	1.0	600	HD 18620
UGPS 0722–05	SpeX/IRTF	2011 Jan 26	LowRes15	0.5	1440	HD 50931
WISEPC J1405 + 5534	WFC3/ <i>HST</i>	2011 Mar 14	G141	...	2212	...
WISEP J1541–2250	FIRE/Magellan	2011 Mar 27	Longslit	0.6	1522	HD 130755
WISEP J1738+2732	WFC3/ <i>HST</i>	2011 May 12	G141	...	2012	...
WISEP J1828+2650	WFC3/ <i>HST</i>	2011 May 9	G141	...	2012	...
WISEPC J2056 + 1459	NIRSPEC/Keck	2010 Oct 21	Low-res (N3)	0.38	2400	HD 198070
	NIRSPEC/Keck	2010 Nov 22	Low-res (N5)	0.38	1800	HD 198069

3.1. Near-infrared Imaging

3.1.1. NEWFIRM/Blanco

WISEP J1541–2250 was observed on the night of 2011 April 17 (UT) with the NOAO Extremely Wide Field Infrared Imager (NEWFIRM) mounted on the Cerro Tololo Inter-American Observatory (CTIO) Victor M. Blanco 4 m Telescope. A description of the instrument, observing strategy, and data reduction can be found in Kirkpatrick et al. (2011). The resultant *J*- and *H*-band photometry is presented in Table 2.

3.1.2. WIRC/Palomar

Near-infrared images of WISEP J0410 + 1502, WISEPC J1405 + 5534, WISEP J1738+2732, and WISEPC J2056 + 1459 were obtained using the Wide-field Infrared Camera (WIRC; Wilson et al. 2003) on the 200 inch Hale Telescope at Palomar Observatory. A description of the instrument, observing strategy, and data reduction can be found in Kirkpatrick et al. (2011). The magnitudes and/or limits for each brown dwarf are given in Table 2.





Figure 1. 2×2 arcmin DSS I, 2MASS *J* and *H*, WISE *W1*, *W2*, and *W3*, and a W1W2W3 false color composite of the five new WISE brown dwarfs. In the color composite images on the far right, the *W1*, *W2*, and *W3* bands are color coded blue, green, and red, respectively.

(A color version of this figure is available in the online journal.)

**Figure 1.** (Continued)

(A color version of this figure is available in the online journal.)

3.1.3. PANIC/Magellan

WISEPC J0148–7202 was observed on the night of 2010 August 1 (UT) with the now decommissioned Persson’s Auxiliary Nasmyth Infrared Camera (PANIC; Martini et al. 2004) on the east Nasmyth platform at the Magellan 6.5 m Baade Telescope. A description of the instrument, observing strategy, and data reduction can be found in Kirkpatrick et al. (2011). The *J*- and *H*-band magnitudes of WISEPC J0148–7202 are given in Table 2.

3.1.4. NIRC2/Keck II

High-resolution observations of WISEP J1828+2650 and WISEPC J2056+1459 were obtained with NIRC2 behind the Keck II LGS-AO system (Wizinowich et al. 2006; van Dam et al. 2006) on the night of 2010 July 1 (UT). A description of the instrument, observing strategy, and data reduction can be found in Kirkpatrick et al. (2011). The *J*- and *H*-band magnitudes are given in Table 2.

3.2. Near-infrared Spectroscopy

3.2.1. SpeX/IRTF

A 0.9–2.5 μm , low-resolution ($R \equiv \lambda/\Delta\lambda \approx 150$) spectrum of UGPS 0722–05 was obtained with SpeX (Rayner et al. 2003) on the 3 m NASA Infrared Telescope Facility (IRTF) on 2011 January 26 (UT). A description of the instrument, observing strategy, and data reduction can be found in Kirkpatrick et al. (2011). The spectrum, which is shown in Figure 2, has a high signal-to-noise ratio (S/N), reaching >50 at the peaks of the *Y*, *J*, and *H* bands.

3.2.2. FIRE/Magellan

Low-resolution ($R = 250\text{--}350$), 1–2.4 μm spectra of WISEPC J0148–7202, WISEP J0410+1502, and WISEP J1541–2250 were obtained with the Folded-port InfraRed Echellette (FIRE; Simcoe et al. 2008, 2010) mounted at the auxiliary Nasmyth focus of the Magellan 6.5 m Baade Telescope. A description of the instrument, observing strategy, and data reduction can be found in Kirkpatrick et al. (2011). The spectra are shown in Figure 2.

3.2.3. NIRSPEC/Keck II

WISEPC J2056+1459 was observed using the Near-Infrared Spectrometer (NIRSPEC; McLean et al. 1998, 2000) located on one of the Nasmyth platforms of the 10 m Keck II telescope on Mauna Kea, Hawaii. The 0′.38-wide slit in the low-resolution mode provides a resolving power of $R = 2500$. WISEPC J2056+1459 was observed with the N3 order sorting filter (1.143–1.375 μm) on the night of 2010 October 21 (UT) and with the N5 order sorting filter (1.431–1.808 μm) on the night of 2010 November 22 (UT).

A series of 300 s exposures was obtained at two different positions along the 42′′ long slit. An A0 V star was observed after each series of science exposures for telluric correction and flux calibration purposes. Calibration frames consisting of neon and argon arc lamps, dark frames, and flat-field lamps were also taken following the science exposures. The data were reduced in a standard fashion using the IDL-based REDSPEC¹⁴ reduction package as described in McLean et al. (2003). Since REDSPEC does not produce uncertainty arrays, we generated them as

follows. First, we performed a simple sum extraction using the rectified, pair-subtracted images generated by REDSPEC. We then scaled the spectra to a common flux level and computed the average spectrum. The uncertainty at each wavelength is given by the standard error on the mean. The average spectrum is then corrected for telluric absorption and flux calibrated using the calibration spectrum generated by REDSPEC. Since the difference between the spectra produced by REDSPEC and our spectra was negligible, we used our spectrum for our analysis. Finally, the N3- and N5-band spectra were absolutely flux calibrated using the WIRC photometry (see Table 2) as described in Cushing et al. (2005) and merged to produce a 1.15–1.80 μm spectrum. The final spectrum, which is shown in Figure 2, has a peak S/N of 8 and 6 in the *J* and *H* bands, respectively.

3.2.4. WFC3/Hubble Space Telescope

WISEPC J1405+5534, WISEP J1738+2732, and WISEP J1828+2650 were observed with the infrared channel of the Wide Field Camera 3 (WFC3; Kimble et al. 2008) on-board the *Hubble Space Telescope* (HST) as a part of a Cycle 18 program (GO-12330, PI: J. D. Kirkpatrick). The WFC3 uses a 1024 × 1024 HgCdTe detector with a plate scale of 0′.13 pixel^{−1} which results in a field of view of 123×126 arcsec. The G141 grism was used to perform slitless spectroscopy of each brown dwarf covering the 1.07–1.70 μm wavelength range at a resolving power of $R \approx 130$. For each brown dwarf, we first obtained four direct images through the F140W filter ($\lambda_p = 1392.3$ nm) in the MULTIACCUM mode with the SPARS25 sampling sequence. Between each exposure, the telescope was offset slightly. We then obtained four images with the G141 grism at the same positions as the direct images. The spectroscopic observations were also obtained in the MULTIACCUM mode but using the SPARS50 sequence.

The raw images were first processed using the CALWFC3 pipeline (ver. 2.3) which not only subtracts the bias level and dark current but also flat fields the direct images (the grism images are flat fielded during the extraction process described below). The spectra were then extracted using the aXe software (Kümmel et al. 2009), which is a suite of PyRAF/IRAF packages designed to extract spectra from the slitless modes of both WFC3 and the Advanced Camera for Surveys (ACS). aXe requires knowledge of both the position and brightness of the objects in the field of view. We therefore combined the four direct images using MULTIDRIZZLE (Koekemoer et al. 2002) and the latest Instrument Distortion Coefficient Table (IDCTAB). A catalog of the objects in the field was then constructed using SExtractor (Bertin & Arnouts 1996). For each object in the source catalog, two-dimensional (2D) subimages centered on the first-order spectra of each object were then combined using the task AXEDRIZZLE to produce a high S/N 2D spectral image. One-dimensional, flux-calibrated spectra and their associated uncertainties are then extracted from the 2D drizzle subimages.

Since the G141 grism mode is slitless, spectral contamination from nearby sources is not uncommon. The aXe software (using the Gaussian emission model) estimates the level of contamination for each object using the positions and magnitudes of all the objects in the field of view. The spectrum of one of the brown dwarfs, WISEP J1828+2650, exhibits moderate contamination that increases in intensity toward shorter wavelengths (see Figure 3). The aXe software does not actually correct for this contamination so we attempted to do

¹⁴ See <http://www2.keck.hawaii.edu/inst/nirspec/redspec>.




Figure 2. Near-infrared spectra of the new *WISE* brown dwarfs (black) as compared to the spectrum of UGPS 0722–05 (red). The data have been normalized to unity at the peak of the *J* band (except for WISEP J1828+2650 which is normalized to unity at the peak of the *H* band) and offset by constants (dotted lines).

(A color version of this figure is available in the online journal.)




Figure 3. Top: subimage of the drizzled WFC3/*HST* grism image centered on the position of WISEP J1828+2650. The location of the spectrum of WISEP J1828+2650 is indicated in red along with the positions of the *J*- and *H*-band peaks. The location of the contaminating starlight is shown in purple and consists of second- and third-order light from two other stars in the WFC3 field of view. Bottom: the spectrum of WISEP J1828+2650 (red) and the contamination spectrum (purple). The stellar contamination becomes progressively worse at shorter wavelengths.

(A color version of this figure is available in the online journal.)

so using the contamination image generated by aXe. Unfortunately, the contamination-corrected spectrum exhibits negative flux values which suggests that aXe is overestimating the contamination level. We will therefore use the contaminated spectrum and consider it an upper limit to the actual spectrum. This issue will be discussed in more detail in Section 4.1.1.

4. ANALYSIS

4.1. Spectral Characterization

Figure 2 shows the near-infrared spectra of the new brown dwarfs. Also plotted for comparison purposes is our IRTF/SpEx spectrum of UGPS 0722–05, the latest-type brown dwarf known previous to this work. All of the spectra exhibit deep H_2O and CH_4 absorption bands characteristic of late-type T dwarfs but the *J*-band peaks of the *WISE* brown dwarfs are narrower than the corresponding peak in the spectrum of UGPS 0722–05. This peak becomes progressively narrower beyond T8 (Warren et al. 2007; Delorme et al. 2008a; Burningham et al. 2008; Lucas et al. 2010), indicating that all of the *WISE* brown dwarfs have spectral types later than UGPS 0722–05. The spectrum of WISEP J1828+2650 is markedly different than that of UGPS 0722–05 so we discuss this object in more detail in the following section before discussing the other six dwarfs.

4.1.1. WISEP J1828+2650: The Archetypal Y Dwarf

The lower panel of Figure 4 shows the 1.15–1.70 μm spectrum of WISEP J1828+2650 along with the spectrum of UGPS 0722–05. The spectrum of WISEP J1828+2650, while dominated by the same CH_4 and H_2O absorption bands present in T dwarf spectra, has a feature not seen in any T dwarf: the *J*- and *H*-band peaks, when plotted in units of f_λ , are essentially




Figure 4. 1.15–1.70 μm spectra of WISEP J1738+2732, WISEPC J1405+5534, and WISEP J1828+2650 along with the spectrum UGPS 0722–05. The uncertainties in the spectra are shown as gray bars. The spectra were all normalized to unity at the peak of the *H* band (1.58 μm). Prominent molecular absorption bands are indicated.

(A color version of this figure is available in the online journal.)

the same height. As discussed in Section 3.2.4, the spectrum of WISEP J1828+2650 is contaminated by light from nearby stars. This contamination, which is *not* removed by the aXe software, becomes progressively worse at shorter wavelengths (see Figure 3) such that the true spectrum will have an even more extreme *J*- to *H*-band peak flux ratio.

The roughly equal-intensity *J* and *H* flux peaks are also confirmed by our ground-based, near-infrared photometry, which gives $J - H = 0.72 \pm 0.42$ mag. Model atmospheres of cool brown dwarfs predict that the near-infrared colors, which are blue for the hotter T dwarfs, turn back to the red at effective temperatures between 300 and 400 K as the Wien tail of the spectral energy distribution collapses. This turn to the red was proposed as one of the triggers that might force the creation of a Y spectral class (Burrows et al. 2003; Kirkpatrick 2008).

Further underscoring the extreme nature of WISEP J1828+2650 is its *J*–*W*2 color of 9.29 ± 0.35 which is over 2 mag redder than the WISEP J1541–2250, the second reddest brown dwarf in our sample at J –*W*2 = 7.18 ± 0.38 (Kirkpatrick et al. 2011). WISEP J1828+2650 is also the reddest brown dwarf in our sample in *H*–*W*2, *J*–[4.5], and *H*–[4.5], where [4.5] represents the *Spitzer Space Telescope* (Werner et al. 2004) Infrared Array Camera (IRAC; Fazio et al. 2004) channel 2 magnitude. Given the extreme nature of both its near-infrared spectrum and near- to mid-infrared colors, we identify WISEP J1828+2650 as the archetypal member of the Y spectral class.




Figure 5. Top: H -band spectrum of 2MASS J22541892+3123498, SDSS J162414.37+002915.6, and 2MASS J04151954-0935066, the T4, T6, and T8 spectral standards (Burgasser et al. 2006), along with the spectrum of UGPS 0722-05, and WISEP J1738+2732. The spectra have been normalized to unity at their peak flux values. Bottom: opacity data computed in chemical equilibrium for NH_3 (Yurchenko et al. 2011), H_2O (Freedman et al. 2008), and CH_4 (Freedman et al. 2008) at $T = 600$ K and $P = 1$ bar. Note that the change in the spectral morphology of the blue wing of the H -band peak is similar between T6/T8 and T8/T9 suggesting a common absorber or set of absorbers. In contrast, the spectrum of WISEP J1738+2732 exhibits excess absorption that matches the position of the NH_3 absorption shown in the lower panel. (A color version of this figure is available in the online journal.)

4.1.2. WISEP J1738+2732 and WISEPC J1405+5534 and the 1.5 μm NH_3 Band

The 1.15–1.70 μm spectra of WISEPC J1405+5534 and WISEP J1738+2732 are very similar and yet both are distinct from UGPS 0722-05, albeit in less extreme ways than the spectrum of WISEP J1828+2650 (see middle and upper panels of Figure 4). Although the relative heights of the J - and H -band peaks are similar to those of UGPS 0722-05, their widths are narrower. The narrowing of the H -band peaks is asymmetric, however, as most of the change is a result of enhanced absorption on the blue wings from 1.51 to 1.58 μm . What is the underlying cause of this absorption?

The H -band spectra of T dwarfs are shaped by CH_4 (and to a lesser extent H_2O) longward of 1.6 μm and by H_2O at wavelengths shortward of 1.6 μm . As the effective temperature falls, the opacity of the near-infrared overtone and combination bands of NH_3 becomes important since $\text{NH}_3/\text{N}_2 > 1$ for $T \lesssim 700$ K at $P = 1$ bar (Lodders & Fegley 2002). The emergence of these NH_3 bands has long been suggested as the trigger for a new spectral class (Burrows et al. 2003; Kirkpatrick 2008; Leggett et al. 2007a) but identifying them has proven difficult because they overlap with the strong H_2O bands and because the abundance of NH_3 can be reduced by an order of magnitude due to vertical mixing in the atmospheres of brown dwarfs (Saumon et al. 2003, 2006; Hubeny & Burrows 2007).

Figure 5 shows the H -band spectra of the T4, T6, and T8 spectral standards, UGPS 0722-05, and WISEP J1738+2732 as well as the opacities for H_2O (Freedman et al. 2008), NH_3 (Yurchenko et al. 2011), and CH_4 (Freedman et al. 2008) at $T = 600$ K and $P = 1$ bar generated by one of us (R.S.F.). With increasing spectral type, the blue wing of the H -band

peak becomes progressively suppressed. Delorme et al. (2008a) tentatively identified NH_3 absorption on the blue wing of the H -band spectrum of CFBDS J0059-0114, a T dwarf with a spectral type earlier than UGPS 0722-05. However, the change in the shape of the blue wing of the H -band peak from T6 to UGPS 0722-05 appears smooth, suggesting a common absorber or set of absorbers. It seems unlikely that NH_3 dominates given that it has not been identified in the spectra of mid-type T dwarf ($T_{\text{eff}} \sim 1200$ K). A similar conclusion to ours is reached by Burningham et al. (2010) using spectral indices.

In contrast, the H -band spectrum of WISEP J1738+2732 stands out in the sequence in that it exhibits additional absorption from 1.53 to 1.58 μm . This absorption broadly matches the position of the $\nu_1 + \nu_3$ absorption band of NH_3 centered at 1.49 μm suggesting that NH_3 is the cause of this absorption. However, we cannot conclusively identify NH_3 as the carrier given the low spectral resolution of the data and the fact that the absorption lies on the steep wing of the H_2O band. For example, water ice also has an absorption band centered at $\sim 1.5 \mu\text{m}$ (Warren & Brandt 2008) that could potentially produce such absorption if the abundance of water ice is high enough. One potential avenue for confirming that NH_3 is indeed the carrier would be to acquire higher spectral resolution data to search for individual NH_3 features (e.g., Saumon et al. 2000; Warren et al. 2007).

4.1.3. The T Dwarf/Y Dwarf Transition

With WISEP J1828+2650 classified as the prototypical Y dwarf, we can now investigate the transition between the T and Y spectral classes. T dwarfs are classified at near-infrared wavelengths using the Burgasser et al. (2006) scheme, wherein nine T dwarf spectral standards with subtypes ranging from T0 to T8 are used for direct spectral comparisons. Burgasser et al. also defined five spectral indices that measure the depths of the CH_4 and H_2O bands which can be used as a proxy for direct comparisons. With the discovery of brown dwarfs with spectral types later than T8, the question of how to extend the Burgasser et al. scheme beyond T8 naturally arises.

The first $>\text{T8}$ dwarf to be identified was ULAS J003402.77-005206.7 (ULAS J0034-0052; Warren et al. 2007). Based on a direct comparison to the spectrum of the T8 spectral standard and the values of the Burgasser et al. spectral indices, Warren et al. adopted a spectral type of T8.5. A second $>\text{T8}$ dwarf soon followed with the discovery of CFBDS J0059-0114 by Delorme et al. (2008a). They used the W_J index which measures the half-width of the J -band peak (Warren et al. 2007) and the NH_3 - H index which measures the half-width of the H -band peak (i.e., the depth of the putative NH_3 absorption), to classify both CFBDS J0059-0114 and ULAS J0034-0052 as T9 dwarfs. Burningham et al. (2008) added two T dwarfs to the tally of $>\text{T8}$ dwarfs with the discovery of ULAS J133553.45+113005.2 (ULAS J1335+1130) and ULAS J123828.51+095351.3 (ULAS J1238+0953). Using both direct spectral comparison and spectral indices, they classified them as T8.5 and T9, respectively. Burningham et al. also proposed extending the Burgasser et al. scheme to T9 by assigning ULAS J1335+1130 as the T9 spectral standard. Additional T dwarfs with spectral types later than T8 have since been discovered (see Table 4), but to date, the latest-type T dwarf currently known is UGPS 0722-05 (Lucas et al. 2010) which has been classified as T10 via a combination of spectral indices and direct comparisons to the T9 dwarfs.




Figure 6. IRTF/SpeX spectra of the Burgasser et al. (2006) spectral standards, SDSS J162414.37 + 002915.6 (T6), 2MASS J07271824 + 1710012 (T7), and 2MASS J04151954 + 0935066 (T8), our IRTF/SpeX spectrum of UGPS 0722–05 and the WFC3/*HST* spectrum of WISEP J1738+2732. The spectra have been normalized to unity at their peak flux level in each panel. Prominent molecular absorption bands are indicated in the top panel.

(A color version of this figure is available in the online journal.)

Figure 6 shows the 1.15–1.70 μm spectra of the T6, T7, T8 spectral standards, UGPS 0722–05, and WISEP J1738+2732. The spectra show smooth changes in their spectral morphology with increasing spectral type including progressively deeper absorption bands centered at 1.15, 1.45, and 1.65 μm and progressively narrower *J*- and *H*-band peaks. However, UGPS 0722–05 does not appear to be two subtypes later than T8 as required by its T10 spectral type. Rather, the changes in spectral morphology from T6 to UGPS 0722–05 suggest that UGPS 0722–05 is more naturally classified as a T9. Given its brightness ($J = 16.5$, 1.5 mag brighter than ULAS J1335 + 1130) and position near the celestial equator, UGPS 0722–05 also makes an ideal spectral standard. We therefore define it to be the T9 spectral standard.

WISEP J1738+2732 is clearly of later type than UGPS 0722–05 but should it be classified as a T dwarf or a Y dwarf? As noted in the previous section, WISEP J1738+2732 exhibits excess absorption from 1.53 to 1.58 μm that we have tentatively ascribed to NH_3 . This absorption becomes even more apparent when the spectrum is placed in sequence with the T6 to T9 spectral standards (lower right panel of Figure 6). Given the smooth change in width of the *J*-band peak and the rapid fall in the flux of the blue wing of the *H* band between UGPS 0722–05 and WISEP J1738+2732 (which suggests the emergence of a new absorption band), we classify WISEP J1738+2732 as a Y dwarf and assign it a spectral type of

Y0. Additionally, we tentatively identify it as Y0 spectral standard.

4.1.4. Classification of the Other WISE Discoveries

With the T9 and Y0 spectral standards defined, we can return to the question of classifying the other new *WISE* discoveries. The *J*- and *H*-band peaks of WISEPC J0148–7202 are slightly narrower than UGPS 0722–05 and slightly wider than WISEP J1738+2732 so we classify this dwarf as T9.5. The spectrum of WISEPC J1405+5534 is very similar to that of WISEP J1738+2732 (see Figure 4). However, we note that the wavelength at which the peak *H*-band flux is reached is shifted ~ 60 Å to the red relative to UGPS 0722–05 (see Figure 4) and the other late-type T dwarfs which suggests that WISEPC J1405+5534 may be peculiar. We therefore classify it as Y0 (pec?). Interestingly, a similar, albeit larger, shift of 200 Å is seen in the spectrum of Jupiter.

The spectra of the remaining brown dwarfs, WISEPC J2056 + 1459, WISEP J0410 + 1502, and WISEP J1541–2250, do not have sufficient S/N to convincingly show the excess absorption from 1.53 to 1.58 μm . However, the *J*-band peaks of these three brown dwarfs are all narrower than UGPS 0722–05. Indeed, the spectra of all of them are a better match to the spectral morphology of WISEP J1738+2732 than UGPS 0722–05 so we classify these brown dwarfs as Y0 as well. In addition, the peak *Y*-band fluxes of WISEP J0410 + 1502 and WISEP




Figure 7. Illustration of the $\text{H}_2\text{O}-J$, W_J , CH_4-J , CH_4-H , NH_3-H , and CH_4-H indices overplotted on the spectrum UGPS 0722-05.

(A color version of this figure is available in the online journal.)

J1541-2250 are slightly higher than in the spectrum of UGPS 0722-05. This is consistent with the blueward trend in the $Y-J$ color of late-type T dwarfs (Leggett et al. 2010; Burningham et al. 2010) which Burningham et al. (2010) ascribed to the weakening of the strong resonance K I doublet (7665, 7699 Å) as K condenses into KCl. Finally, since the spectrum of WISEP J1828+2650 is distinct from both UGPS 0722-05 and WISEP J1738+2732, we classify it as $>Y0$. A more precise subtype will require the discovery of additional Y dwarfs to bridge the gap in spectral morphology between WISEP J1738+2732 and WISEP J1828+2650.

4.1.5. Reclassification of Previously Published $\geq T8.5$ Dwarfs

There are also 12 T dwarfs with spectral types later than T8 currently in the literature (see Table 4). Since we have reclassified UGPS 0722-05 as a T9 dwarf, we must also reclassify the other 11 dwarfs using this new system. To accomplish this, we have smoothed the published spectra to a resolving power of $R = 150$ and resampled them onto the same wavelength scale as the IRTF/SpeX spectrum of UGPS 0722-05. This ensures that differences in resolving power and wavelength sampling between the late-type T dwarfs and the Burgasser et al. IRTF/SpeX spectra of the T dwarf spectral standards do not adversely affect our classification. Table 4 gives the revised spectral types derived from direct comparison for the 12 T dwarfs with published spectral types later than T8. When the J - and H -band regions gave conflicting spectral types, we used the typed inferred from the J band.

4.1.6. Spectral Indices

Although the primary (and preferred) method of assigning a spectral type is to compare the spectrum of a brown dwarf against that of the spectral standards, the use of spectral indices remains popular in the literature. We have therefore computed the $\text{H}_2\text{O}-J$, CH_4-J , $\text{H}_2\text{O}-H$, CH_4-H (Burgasser et al. 2006), W_J (Warren et al. 2007), and NH_3-H (Delorme et al. 2008a) indices of the new WISE brown dwarfs and as well as UGPS 0722-05. Figure 7 illustrates the positions of the indices'

flux windows relative to the spectrum of UGPS 0722-05. The index values are computed in a Monte Carlo fashion whereby 5000 realizations of each spectrum are generated by randomly drawing from normal distributions with means given by the flux densities at each wavelength and standard deviations given by the uncertainty in the flux densities. The values of the indices and their uncertainties are given by the mean and standard deviation of the distribution of index values computed from the 5000 realizations and are listed in Table 5.

Figure 8 shows the values of the six indices as a function of spectral type. Also shown are the index values of the T6-T8 spectral standards, a sample of T5-T8 dwarfs from the SpeX Prism Spectral Library, and the 12 T dwarfs with previously published spectral types later than T8. The classification of WISEP J1738+2732 as a Y dwarf is bolstered by the distinctive break in the trend of the NH_3-H values with spectral type which suggests that a new absorption band has indeed emerged at the T/Y dwarf transition. The remaining spectral indices do not show a break at the T/Y transition, but the CH_4-J , $\text{H}_2\text{O}-H$, and W_J indices do show a smooth trend with spectral type down to Y0 indicating that they can still be used as proxies for direct spectral comparisons. Indeed, the value of the W_J spectral index for WISEP J1738+2732 is far from saturated so we support the suggestion by Burningham et al. (2008) that this index can be used as a proxy for direct comparison for late-type T dwarfs and early-type Y dwarfs. However, the CH_4-H index is clearly beginning to saturate at T9 and the $\text{H}_2\text{O}-J$ index may even reverse at Y0 rendering these indices less useful for classification purposes. Finally, although there is scatter due to the very low S/N of some of the spectra, the new WISE brown dwarfs are clearly distinct from the previous $\geq T8.5$ dwarfs and cluster around the Y0 spectral standard.

4.2. Atmospheric and Structural Properties

4.2.1. Atmospheric Properties

In order to investigate the atmospheric properties (e.g., T_{eff} , $\log g$) of the brown dwarfs, we have compared their near-infrared spectra to a new preliminary grid of model spectra generated with the model atmospheres of Marley & Saumon. A detailed description of the basic models can be found in Marley et al. (2002), Saumon & Marley (2008), Cushing et al. (2008), and Stephens et al. (2009). This preliminary grid includes a new NH_3 line list (Yurchenko et al. 2011) and a new prescription for the collision induced opacity for H_2 (D. Saumon et al. 2011, in preparation). A more detailed study that compares the model spectra to the near-infrared spectra, and WISE and *Spitzer* photometry is in preparation.

The grid consists of solar metallicity, cloudless models with the following parameters: $T_{\text{eff}} = 200\text{--}1000$ K in steps of 50 K, $\log g = 3.75 - 5$ in steps of 0.25 (cm s^{-2}), and $K_{\text{zz}} = 0, 10^4 \text{ cm}^2 \text{ s}^{-1}$. Although the opacities of the condensate clouds are not included in the atmospheric models, i.e., they are cloudless, the effects on the atmospheric chemistry due to the rainout of the condensates is accounted for in the models. This assumption is reasonable for the silicate and liquid iron clouds since they form well below the observable photosphere (see however, Burgasser et al. 2010) but may not be valid if, as expected, H_2O clouds form high in the atmosphere of the coldest models. The eddy diffusion coefficient, K_{zz} , parameterizes the vigor of mixing in the radiative layers of the atmosphere. A value of $K_{\text{zz}} > 0 \text{ cm}^2 \text{ s}^{-1}$




Figure 8. Values of the $\text{H}_2\text{O}-J$, CH_4-J , $\text{H}_2\text{O}-H$, CH_4-H (Burgasser et al. 2006), W_J (Warren et al. 2007), and NH_3-H (Delorme et al. 2008a) spectral indices as a function of spectral type. The black points are for the T6–Y0 spectral standards. The gray points were computed using spectra of late-type (T5–T8) T dwarfs from the SpeX Prism Spectral Library. The red points are the 12 T dwarfs with previously published spectral types later than T8 and the green points are the remaining six *WISE* brown dwarfs. For plotting purposes only, we have assigned WISEP J1828+2650 a spectral type of Y1.

(A color version of this figure is available in the online journal.)

Table 4
Previously Published Brown Dwarfs with Spectral Types Later than T8

Object	Previous Spectral Type	Reference	Adopted Spectral Type
Ross 458C	T8	Burgasser et al. (2010)	T8
	T8.5p	Burningham et al. (2011a)	...
ULAS J123828.51 + 095351.3	T8.5	Burningham et al. (2008)	T8
ULAS J130217.21 + 130851.2	T8.5	Burningham et al. (2010)	T8
ULAS J003402.77–005206.7	T8.5	Warren et al. (2007)	T8.5
	T9	Burningham et al. (2008)	...
CFBDS J005910.90–011401.3	\gtrsim T9	Delorme et al. (2008a)	T8.5
	T9	Burningham et al. (2008)	...
WISEPC J045853.90 + 643451.9	T9	Mainzer et al. (2011)	T8.5
UGPS J052127.27 + 364048.6	T8.5	Burningham et al. (2011b)	T8.5
ULAS J133553.45 + 113005.2	T9	Burningham et al. (2008)	T8.5
Wolf 940B	T8.5	Burningham et al. (2009)	T8.5
WISEPC J181210.85 + 272144.3	T8.5:	Burgasser et al. (2011)	T8.5:
UGPS J072227.51–054031.2	T10	Lucas et al. (2010)	T9
CFBDSIR J145829 + 101343AB	T9.5	Liu et al. (2011)	T9

results in mixing that can prevent the abundances of CO and CH_4 (the dominant carbon-bearing species) from coming into chemical equilibrium because the mixing timescales become shorter than the timescales of key chemical reactions involved in the conversion of CO to CH_4 (Lodders & Fegley 2002; Saumon et al. 2003; Hubeny & Burrows 2007). Typical values of K_{zz} in the stratospheres of giant planets are 10^2 – $10^5 \text{ cm}^2 \text{ s}^{-1}$ (Saumon et al. 2006). The abundances of N_2 and NH_3 (the

dominant nitrogen-bearing species) are also kept from coming into chemical equilibrium by mixing, but in this case the mixing timescales are set in the convective layers of the atmosphere by the mixing length theory. As a result, the final non-equilibrium abundances of N_2 and NH_3 are not sensitive to variations in the eddy diffusion coefficient K_{zz} . However by convention, models with $K_{zz} = 0 \text{ cm}^2 \text{ s}^{-1}$ are in full chemical equilibrium (i.e., the effect of convective mixing on the nitrogen chemistry is

Table 5
Spectral Indices

Object	Spectral Type	H ₂ O- <i>J</i>	CH ₄ - <i>J</i>	H ₂ O- <i>H</i>	CH ₄ - <i>H</i>	<i>W_J</i>	NH ₃ - <i>H</i>
UGPS 0722-05 ^a	T9	+0.009 (0.004)	+0.115 (0.003)	+0.115 (0.007)	+0.075 (0.005)	+0.215 (0.003)	+0.527 (0.008)
WISEPC J0148-7202	T9.5	-0.017 (0.011)	+0.076 (0.006)	+0.070 (0.013)	+0.016 (0.012)	+0.152 (0.006)	+0.431 (0.015)
WISEP J0410+1502	Y0	-0.043 (0.090)	+0.036 (0.059)	-0.036 (0.458)	-0.105 (0.434)	+0.155 (0.052)	+0.380 (0.574)
WISEPC J1405+5534	Y0 (pec?)	+0.020 (0.026)	+0.031 (0.017)	+0.063 (0.025)	+0.162 (0.028)	+0.099 (0.017)	+0.346 (0.027)
WISEP J1541-2250	Y0	-0.046 (0.115)	+0.040 (0.060)	+0.155 (0.216)	+0.057 (0.208)	+0.081 (0.066)	+0.259 (0.200)
WISEP J1738+2732	Y0	+0.036 (0.008)	+0.050 (0.005)	+0.045 (0.008)	+0.050 (0.009)	+0.149 (0.005)	+0.349 (0.010)
WISEP J1828+2650	>Y0	+0.248 (0.152)	+0.257 (0.099)	+0.049 (0.091)	+0.129 (0.101)	+0.515 (0.106)	+0.503 (0.110)
WISEPC J2056+1459	Y0	...	+0.030 (0.009)	...	+0.050 (0.034)	+0.129 (0.012)	+0.240 (0.031)

Notes. The H₂O-*J* and H₂O-*H* indices cannot be computed for WISEPC J205628.90+145953.3 because its spectrum does not span the entire wavelength range of the indices.

^a The values differ from that measured by Lucas et al. (2010). Our two spectra agree well except deep in the CH₄ and H₂O absorption bands, where our spectrum exhibits lower flux levels. The reason for this discrepancy is unclear but it may be a result of the fact that the Lucas et al. spectrum was created by merging separate spectra that were absolutely flux calibrated using near-infrared photometry.

ignored) and models with $K_{zz} \neq 0 \text{ cm}^2 \text{ s}^{-1}$ exhibit both carbon and nitrogen non-equilibrium chemistry.

The best-fitting models are identified using the goodness-of-fit statistic, $G_k = \sum_{i=1}^n w_i \left(\frac{f_i - C_k F_{k,i}}{\sigma_i} \right)^2$, where n is the number of data pixels, w_i is the weight for the i th wavelength (set to unity in this case), f_i and $F_{k,i}$ are the flux densities of the data and model k , respectively, σ_i are the errors in the observed flux densities, and C_k is an unknown multiplicative constant equal to $(R/d)^2$, where R is the radius of the star and d is the distance to the star (Cushing et al. 2008). In order to increase the S/N of the data, we first smoothed the higher resolution spectra to $R = 200$. The model spectra were also smoothed to the same resolving power and linearly interpolated onto the wavelength scale of the data. For each model, we compute the scale factor C_k by minimizing G_k with respect to C_k and identify the best-fitting model as having the global minimum G_k value. To estimate the range of models that fits the data well, we run a Monte Carlo simulation that uses the uncertainties in the individual spectral points and the uncertainties in the absolute flux calibration of the spectra to generate 10^4 simulated noisy spectra. The fitting process is repeated on each simulated spectrum and models that are consistent with the best-fitting model at the 3σ level are considered equally good representations of the data. We did not attempt to fit the spectrum of WISEP J1828+2650 because it is contaminated with light from other stars in the WFC3 field of view (see Sections 3.2.4 and 4.1.1). After discussing the results of the fits to the spectra of the other brown dwarfs, we return to estimate an approximate effective temperature for WISEP J1828+2650.

Table 6 lists the best-fitting model parameters for each brown dwarf, as well as UGPS 0722-05. The derived effective temperatures of the *WISE* brown dwarfs are all cold, ranging from 350 to 500 K. Indeed all but one have estimated effective temperatures of less than 450 K making them the coldest spectroscopically confirmed brown dwarfs known. Five out of the six best-fitting models also have $K_{zz} \neq 0$ which indicates that vertical mixing is present in the atmospheres of these cold brown dwarfs. This is not a surprising result given that strong evidence for vertical mixing in the atmospheres of brown dwarfs has been found at longer wavelengths (Saumon et al. 2006, 2007; Leggett et al. 2007b, 2009; Burgasser et al. 2008; Stephens et al. 2009; Geballe et al. 2009; Cushing et al. 2010). At such low temperatures, the effects of non-equilibrium chemistry on the *J*- and *H*-band spectra of brown dwarfs is limited to




Figure 9. Best-fitting models (red) overplotted on the near-infrared spectra of six of the seven new *WISE* brown dwarfs (black). The spectra were normalized to unity at the peak flux in the *J* band and offset by constants (dotted lines). The uncertainties in the spectra are given by gray bars. The best-fitting model parameters are given in the form $T_{\text{eff}}(\text{K})/\log g (\text{cm s}^{-2})/K_{zz} (\text{cm}^2 \text{ s}^{-1})$.

(A color version of this figure is available in the online journal.)

weakening the NH₃ absorption bands. The detection of mixing therefore underscores the fact that NH₃ is probably at least partially responsible for the absorption seen on the blue wing of the *H*-band peak of WISEPC J1405+5534 and WISEP J1738+2732.

Figure 9 shows the best-fitting model spectra overplotted on the data. Since this is the first time such cold model spectra have been compared to observed spectra, the agreement between the models and the data is encouraging. In particular, the height

Table 6
Atmospheric and Structural Properties

Object	SpType	T_{eff} (K)	$\log g$ (cm s^{-2})	$\log K_{zz}$ ($\text{cm}^2 \text{s}^{-1}$)	R (R_{Jup})	M (M_{Jup})
UGPS 0722–05	T9	650	4.00 (4.00–4.25)	4	1.21 (1.14–1.21)	6 (6 – 9)
WISEPC J0148–7202	T9.5	500 (500–500)	4.50 (4.50–4.75)	4	1.04 (0.96–1.04)	13 (13–21)
WISEP J0410 + 1502	Y0	450 (400–500)	3.75 (3.75–4.25)	0	1.22 (1.09–1.22)	3 (3–9)
WISEPC J1405 + 5534	Y0 (pec?)	350	5.00	4	0.86	30
WISEP J1541–2250	Y0	350	4.50 (4.25–4.5)	4	1.01 (1.01–1.07)	12 (8–12)
WISEP J1738+2732	Y0	350 (350–400)	4.75 (4.75–5.00)	4	0.93 (0.86–0.94)	20 (20–30)
WISEP J1828+2650	>Y0	≤ 300
WISEPC J2056 + 1459	Y0	350 (350–400)	4.75 (4.50–5.00)	4	0.93 (0.86–1.01)	20 (12–30)

Notes. The parameters for the best-fitting Marley & Saumon models are listed and the range of parameters consistent with the data is given in parentheses. The effective temperature limit for WISEP J1828+2650 was estimated by identifying those models with peak J -band fluxes equal to or less than the peak flux in the H band and by comparing the observed J – $W2$ color to model J – $W2$ colors.

and width of the J -band peaks are well matched by the model spectra. Previous studies of hotter brown dwarfs fail to match both the peak and width of this peak (e.g., Leggett et al. 2009; Burgasser et al. 2011). The improved fits may be a result of the fact that we are fitting a smaller wavelength range (Cushing et al. 2008; Seifahrt et al. 2009) and/or because the high J vibration–rotation lines (the so-called hot lines), whose cross sections are less well known, become less important at such cold temperatures.

The models do, however, provide a poor fit to the blue wing of the H -band peak of the spectra and fail to reproduce the heights of the Y -band peaks of WISEP J0410 + 1502 and WISEP J1541–2250. Note that the peak of the Y band is shaped by the $2\nu_1+2\nu_4$ band of NH_3 centered at about $1.03 \mu\text{m}$ and therefore Y -band spectra of cold brown dwarfs could provide the first clear detection of NH_3 at near-infrared wavelengths. In principle, the blue wing of the H -band model spectrum could be brought into better agreement with the data by further reducing the abundance of NH_3 . However, as noted above, the abundance of NH_3 is insensitive to variations in K_{zz} because it is quenched in the convective region where the mixing timescale is set by the mixing length theory and not by the eddy diffusion coefficient. Therefore, the mismatch between the data and models is most likely a result of some other inadequacy in the model atmospheres.

Finally, although we cannot fit the models to the spectrum of WISEP J1828+2650, we can still estimate a rough effective temperature. The most salient feature of the spectrum is that the J - and H -band peaks are roughly the same height in flux density units of f_λ . Only model spectra with $T_{\text{eff}} \leq 250$ K have J -band peak fluxes that are equal to or less than the H -band peak fluxes. A second estimate of the effective temperature can be derived from the observed J – $W2$ color of 9.29 ± 0.35 . We computed synthetic Mauna Kea Observatories Near-Infrared (MKO-NIR; Tokunaga et al. 2002) J and $W2$ magnitudes for each model in the grid and find that model spectra with $T_{\text{eff}} = 275$ – 300 K have J – $W2$ colors that fall within $\pm 2\sigma$ of the observed color. Taken together, these estimates suggest that an appropriate upper limit to the effective temperature of WISEP J1828+2650 is ~ 300 K which makes WISEP J1828+2650 the coolest spectroscopically confirmed brown dwarf known.

4.2.2. Structural Properties

With estimates of the effective temperatures and surface gravities of the new brown dwarfs in hand, we can also estimate their radii (R) and masses (M) using evolutionary models. We

Table 7
Distance Estimates

Object	SpType	d_{spec} (pc) ^a	d_π (pc) ^b	d_{phot} (pc) ^c
UGPS 0722–05	T9	11.1 (10.4–11.1)	3.6–4.7	...
WISEPC J0148–7202	T9.5	14.7 (13.1–14.7)	...	12.1
WISEP J0410 + 1502	Y0	11.8 (6.3–16.9)	...	9.0
WISEPC J1405 + 5534	Y0 (pec?)	3.8	...	8.6
WISEP J1541–2250	Y0	8.1 (8.1–8.9)	2.2–4.1	8.2
WISEP J1738+2732	Y0	3.4 (3.4–7.3)	...	10.5
WISEP J1828+2650	>Y0	<9.4
WISEPC J2056 + 1459	Y0	3.0 (2.4–6.4)	...	7.7

Notes.

^a Spectroscopic distance estimates derived as described in Section 4.2.1. The distance corresponding to the best-fitting model is given and the range of distances corresponding to models that are consistent with the data are given in parentheses.

^b Parallaxic distance for UGPS 0722–05 and WISEP J1541–2250 from Lucas et al. (2010) and Kirkpatrick et al. (2011), respectively.

^c Photometric distance estimates from Kirkpatrick et al. (2011).

used the cloudless structure models of Saumon & Marley (2008) because they used atmospheric models that are nearly identical to the ones we used in our analysis for boundary conditions. As a result, the derived T_{eff} , $\log g$, R , and M estimates are all self-consistent. The radii and masses of the brown dwarfs are given in Table 6.

4.3. Spectroscopic Distance Estimates

The value of the multiplicative constant $C_k = (R/d)^2$ derived as a byproduct of the atmospheric model fitting procedure can be used to estimate the so-called spectroscopic distance (d_{spec}) to brown dwarfs if their radii can be determined (e.g., Bowler et al. 2009). In the absence of direct measurements of brown dwarf radii, we can use the radii computed using evolutionary models and (T_{eff} , $\log g$) values. The spectroscopic distances of the new *WISE* brown dwarfs and UGPS 0722–05 are given in Table 7. Also listed in Table 7 are the photometric distance estimates of the *WISE* brown dwarfs from Kirkpatrick et al. (2011) and parallaxic distances to UGPS 0722–05 (Lucas et al. 2010) and WISEP J1541–2250 (Kirkpatrick et al. 2011). The former distance estimates are computed using a spectral type– $W2$ relation derived from known brown dwarfs with spectral types ranging from L0 to T9 and with $\pi/\sigma_\pi > 3$. The photometric distances of the new *WISE* brown dwarfs are based on an




Figure 10. Impact of systematic errors in the derived values of (T_{eff} , $\log g$) on the spectroscopic distance, d_{spec} , for a hypothetical dwarf with $T_{\text{eff}} = 600$ K and $\log g = 4.5 \text{ cm s}^{-2}$.

(A color version of this figure is available in the online journal.)

extrapolation of this relation and therefore should be treated with caution.

The agreement between the three distance estimates range from good to poor. For example, the spectroscopic and photometric distances of WISEP J1541–2250 are in good agreement but the parallaxic distance is discrepant by a factor of two to four. Perhaps most discouraging is the mismatch between the spectroscopic and parallaxic distances of UGPS 0722–05. Liu et al. (2011) recently showed that the agreement between the spectroscopic distances (derived using only near-infrared spectra) and the parallaxic distances of 10 late-type T dwarfs range from 10% to a factor of two, with no apparent trend with spectral type or distance. This suggests that spectroscopic distances should only be used to confirm that the physical properties of brown dwarfs (T_{eff} , R , M) derived from atmospheric and evolutionary models are consistent with the known distance to the brown dwarf.

A corollary to this statement is that if the spectroscopic and parallaxic distances are discrepant then some combinations of the T_{eff} , $\log g$, and R values are in error. In order to estimate the significance of the bias in the spectroscopic distance estimate introduced by systematic errors in the inferred atmospheric properties, we have run a $1\text{--}2.5 \mu\text{m}$ model with $T_{\text{eff}} = 600$ K, $\log g = 4.5 \text{ cm s}^{-2}$ through the fitting procedure described in Section 4.2.1. The model spectrum was first multiplied by an appropriate value of $(R/d)^2$ corresponding to 10 pc. Figure 10 shows the ratio of $d_{\text{spec}}/10 \text{ pc}$ derived for models with effective temperatures from 500 to 700 K and surface gravities from 3.75 to 5.0. The maximum change in d_{spec} for these models is approximately a factor of two for a change of $+100 \text{ K}/-0.75 \text{ dex}$ and $-100 \text{ K}/+0.5 \text{ dex}$ in $T_{\text{eff}}/\log g$. The spectroscopic distance is also most sensitive to changes in T_{eff} as noted by Liu et al. (2011).

The apparent mismatch between the spectroscopic and parallaxic distance estimates is perhaps not so surprising as Cushing et al. (2008) have shown that variations of order 100–200 K are common when estimating the effective temperatures of L and early- to mid-type T dwarfs using spectra that cover only a fraction of the spectral energy distribution. These variations are most likely exacerbated by the fact that only $\sim 35\%$ (in flux density units of f_{λ}) of the bolometric flux of a $T_{\text{eff}} = 600$ K brown dwarf is emitted at near-infrared wavelengths. In summary, given the

uncertainties in the model atmospheres and the difficulty in estimating the effective temperatures and surface gravities of brown dwarfs, it is not surprising that spectroscopic distance estimates do not always agree with the parallaxic distances.

5. DISCUSSION

The new *WISE* brown dwarfs presented herein are the coldest ($T_{\text{eff}} = 300\text{--}500$ K) spectroscopically confirmed brown dwarfs currently known. However as noted in Section 1, WD 0806–661B and CFBDSIR J1458 + 1013B have estimated effective temperatures of $\sim 300\text{--}400$ K based on photometry alone (Luhman et al. 2011; Liu et al. 2011; Rodriguez et al. 2011). How do the properties of these two brown dwarfs compare with the new *WISE* brown dwarfs?

The upper panel of Figure 11 shows the absolute *J*-band magnitude (M_J) as a function of spectral type for a sample of field T dwarfs (Leggett et al. 2010), WISEP J1541–2250 (the only *WISE* brown dwarf with a measured parallax), WD 0806–661B (Luhman et al. 2011) and CFBDSIR J1458 + 1013B (Liu et al. 2011). The value of M_J increases precipitously beyond a spectral type of T8 and reaches ~ 23.9 for WISEP J1541–2250 (Y0). The absolute magnitude of CFBDSIR J1458 + 1013B falls between the two T9 dwarfs and WISEP J1541–2250 suggesting that it has a spectral type of T9–Y0. However, the absolute magnitudes of more late-type T dwarfs and Y dwarfs must be measured before any strong conclusions can be drawn based on absolute magnitudes alone. The absolute magnitude of WD 0806–661B is still only a limit ($M_J > 22.5$) which leaves open the possibility that WD 0806–661B is even fainter, and thus presumably cooler than, WISEP J1541–2250. Either way, it is clear that based on the trend of M_J with spectral type that observing even colder objects at near-infrared wavelengths is going to become increasingly difficult unless they are very close the Sun.

The lower panel of Figure 11 shows the *J*–*H* colors as a function of spectral type for a sample of field T dwarfs (Leggett et al. 2010), the new *WISE* brown dwarfs, and CFBDSIR J1458 + 1013B (WD 0806–661B has yet to be detected in either the *J* or *H* band). Some of the colors of the *WISE* brown dwarfs have large uncertainties so we also computed synthetic MKO-NIR *J*–*H* colors as described in Rayner et al. (2009); they are shown as triangles in Figure 11. Since the WFC3/*HST* spectra do not span the entire *H* band, we used the spectrum of UGPS 0722–05 to extend the spectra of WISEP J1738+2732 and WISEPC J1405+5534 to the limit of the *H*-band filter. The synthetic and observed photometries of WISEPC J1405 + 5534 are clearly discrepant and it is unclear what the underlying cause is. The scatter in the *J*–*H* colors of the dwarfs at the T/Y transition makes it difficult to assign a spectral type to CFBDSIR J1458 + 1013B but it is broadly consistent with T6–Y0. The overall trend with spectral type suggests that the *J*–*H* colors may be turning toward the red at the T/Y transition. This turn is broadly consistent with theoretical models which predict that the *J*–*K* color also turns toward the red at $T_{\text{eff}} = 390\text{--}450$ K (Burrows et al. 2003). However, given the small number of objects and the large uncertainties in colors, a definitive conclusion cannot yet be reached.

Finally, given the rapid increase in the absolute *J*-band magnitude at the T/Y transition, it is reasonable to ask whether the near-infrared is the appropriate wavelength range with which to define the Y spectral class. Indeed historically as cooler and cooler stars were discovered, the wavelength range used for spectral classification moved from the blue violet at $3900\text{--}4900 \text{ \AA}$ (Morgan et al. 1943), through the red optical




Figure 11. Comparison of the absolute J -band magnitudes and $J-H$ colors of the *WISE* brown dwarfs, WD 0806–661B, and CFBDSIR J1458 + 1013B. The field population (black circles) is from the compilation of Leggett et al. (2010) and the spectral types of the late-type T dwarfs have been changed to match the subtypes given in Table 4. Synthetic colors computed by integrating the J and H bandpasses over spectra are plotted as triangles. For plotting purposes only, we have assigned a spectral type of Y1 for WISEP J1828+2650.

(A color version of this figure is available in the online journal.)

at 7000–10000 Å (Boeshaar 1976; Kirkpatrick et al. 1991, 1999), and finally into the near-infrared at 1–2.5 μ m (Burgasser et al. 2006). Since Y dwarfs emit the majority of their flux in the mid-infrared, it seems only natural to devise the spectral classification system at these wavelengths. The smooth spectral morphological changes seen in the 5.5–14.5 μ m *Spitzer* spectra of L and T dwarfs (Cushing et al. 2006) suggest that a mid-infrared classification scheme for the Y dwarfs is plausible. Unfortunately, observing at wavelengths longer about 2.5 μ m from the ground is exceedingly difficult due to the high thermal background. Observations conducted from space do not suffer from this limitation but there are currently no space-based facilities (including *Spitzer*) capable of obtaining mid-infrared spectra of cold brown dwarfs. The *James Webb Space Telescope* will provide a platform with which to observe cold brown dwarfs (Burrows et al. 2003; Marley & Leggett 2009) but its launch is, at best, still years away.

We are therefore left in the unfortunate position of either waiting for a (space- or ground-based) facility capable of sensitive mid-infrared observations or constructing a classification scheme in the near-infrared. Given the large number of cold brown dwarfs now known, we believe it is important to devise a system with which to classify them based on the data currently available. The creation of a near-infrared scheme in no way invalidates any future mid-infrared system that may be de-

vised. Indeed, the classification of brown dwarfs at two different wavelengths is not unprecedented as both the L and the T dwarfs have classification systems based in the red optical (Kirkpatrick et al. 1999; Burgasser et al. 2003) and the near-infrared (Geballe et al. 2002; Kirkpatrick et al. 2010; Burgasser et al. 2006). Ultimately, the utility of any classification system will be measured by whether or not it is adopted by the brown dwarf community.

6. SUMMARY

As part of an ongoing search for the coldest brown dwarfs in the solar neighborhood using the *WISE*, we have discovered seven ultracool brown dwarfs whose near-infrared spectra indicate that they are the latest-type brown dwarfs currently known. Based on the spectral morphological differences between these brown dwarfs and the late-type T dwarfs, we have identified six of them as the first members of the Y spectral class. A comparison to the model spectra of Marley & Saumon indicates that they have effective temperatures ranging from 300 to 500 K and thus are the coolest spectroscopically confirmed brown dwarfs currently known.

We thank Tom Jarrett for guidance with the WIRC data reduction, Barry Rothberg, and Norbert Pirzkal for their guidance in reducing the *HST*/WFC3 data, and Ben Burningham, Sandy Leggett, and Mike Liu for providing digital copies of late-type T dwarf spectra. We also thank Mauricio Martinez, Jorge Araya, and Nidia Morrell for observing support at Magellan. M.S.M. and D.S. acknowledge the support of the NASA ATP program. This publication makes use of data products from the *Wide-field Infrared Survey Explorer*, the Two Micron All Sky Survey (2MASS), and the Sloan Digital Sky Survey (SDSS). The *Wide-field Infrared Survey Explorer* is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration. 2MASS is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. SDSS is funded by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. This research has made use of the NASA/IPAC Infrared Science Archive (IRSA), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Our research has also benefited from the M, L, and T dwarf compendium housed at DwarfArchives.org whose server was funded by a NASA Small Research Grant, administered by the American Astronomical Society and the SpeX Prism Spectral Libraries, maintained by Adam Burgasser at <http://www.browndwarfs.org/speXprism>. Data presented herein were obtained at the W. M. Keck Observatory from telescope time allocated to the National Aeronautics and Space Administration through the agency’s scientific partnership with the California Institute of Technology and the University of California. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation. This paper also includes data gathered with the 6.5 m Magellan Telescopes located at Las Campanas Observatory, Chile and the Peters Automated Infrared Imaging Telescope (PAIRITEL) which is operated by the

Smithsonian Astrophysical Observatory (SAO) and was made possible by a grant from the Harvard University Milton Fund, the camera loan from the University of Virginia, and the continued support of the SAO and the University of California, Berkeley. Magellan telescope time was granted by NOAO (Proposal ID 2010B-0184, P.I. Mainzer), through the Telescope System Instrumentation Program (TSIP). TSIP is funded by NSF National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA) under cooperative agreement with the National Science Foundation. Partial support for PAIRITEL operations and this work comes from National Aeronautics and Space Administration grant NNG06GH50G. A.J.B. acknowledges support from the Chris and Warren Hellman Fellowship Program. Finally, this research was supported (in part) by an appointment to the NASA Postdoctoral Program at the Jet Propulsion Laboratory, administered by Oak Ridge Associated Universities through a contract with NASA.

Facilities: IRTF (SpeX), Hale (WIRC), Magellan:Baade (FIRE, PANIC), Keck:II (NIRC2, NIRSPEC), *Spitzer* (IRAC), *HST* (WFC3)

REFERENCES

- Allen, P. R., Koerner, D. W., Reid, I. N., & Trilling, D. E. 2005, *ApJ*, **625**, 385
- Bertin, E., & Arnouts, S. 1996, *A&AS*, **117**, 393
- Beuzit, J. L., Mouillet, D., Moutou, C., et al. 2006, in Tenth Anniversary of 51 Peg-b: Status of and Prospects for Hot Jupiter studies, ed. L. Arnold, F. Bouchy, & C. Moutou (Paris: Frontier Group), 353
- Boeshaar, P. C. 1976, PhD thesis, Ohio State Univ., Columbus
- Bowler, B. P., Liu, M. C., & Cushing, M. C. 2009, *ApJ*, **706**, 1114
- Burgasser, A. J. 2004, *ApJS*, **155**, 191
- Burgasser, A. J., Cushing, M. C., Kirkpatrick, J. D., et al. 2011, *ApJ*, **735**, 116
- Burgasser, A. J., Geballe, T. R., Leggett, S. K., Kirkpatrick, J. D., & Golimowski, D. A. 2006, *ApJ*, **637**, 1067
- Burgasser, A. J., Kirkpatrick, J. D., Liebert, J., & Burrows, A. 2003, *ApJ*, **594**, 510
- Burgasser, A. J., Simcoe, R. A., Bochanski, J. J., et al. 2010, *ApJ*, **725**, 1405
- Burgasser, A. J., Tinney, C. G., Cushing, M. C., et al. 2008, *ApJ*, **689**, L53
- Burningham, B., Leggett, S. K., Homeier, D., et al. 2011a, *MNRAS*, **414**, L390
- Burningham, B., Lucas, P. W., Leggett, S. K., et al. 2011b, *MNRAS*, **414**, L90
- Burningham, B., Pinfield, D. J., Leggett, S. K., et al. 2008, *MNRAS*, **391**, 320
- Burningham, B., Pinfield, D. J., Leggett, S. K., et al. 2009, *MNRAS*, **395**, 1237
- Burningham, B., Pinfield, D. J., Lucas, P. W., et al. 2010, *MNRAS*, **406**, 1885
- Burrows, A., & Sharp, C. M. 1999, *ApJ*, **512**, 843
- Burrows, A., Sudarsky, D., & Lunine, J. I. 2003, *ApJ*, **596**, 587
- Chauvin, G., Lagrange, A., Dumas, C., et al. 2004, *A&A*, **425**, L29
- Cushing, M. C., Marley, M. S., Saumon, D., et al. 2008, *ApJ*, **678**, 1372
- Cushing, M. C., Rayner, J. T., & Vacca, W. D. 2005, *ApJ*, **623**, 1115
- Cushing, M. C., Roellig, T. L., Marley, M. S., et al. 2006, *ApJ*, **648**, 614
- Cushing, M. C., Saumon, D., & Marley, M. S. 2010, *AJ*, **140**, 1428
- Delorme, P., Delfosse, X., Albert, L., et al. 2008a, *A&A*, **482**, 961
- Delorme, P., Willott, C. J., Forveille, T., et al. 2008b, *A&A*, **484**, 469
- Epchtein, N., de Batz, B., Capozzi, L., et al. 1997, *Messenger*, **87**, 27
- Fazio, G. G., Hora, J. L., Allen, L. E., et al. 2004, *ApJS*, **154**, 10
- Freedman, R. S., Marley, M. S., & Lodders, K. 2008, *ApJS*, **174**, 504
- Geballe, T. R., Knapp, G. R., Leggett, S. K., et al. 2002, *ApJ*, **564**, 466
- Geballe, T. R., Saumon, D., Golimowski, D. A., et al. 2009, *ApJ*, **695**, 844
- Hanel, R., Conrath, B., Herath, L., Kunde, V., & Pirraglia, J. 1981, *J. Geophys. Res.*, **86**, 8705
- Hayashi, C., & Nakano, T. 1963, *Prog. Theor. Phys.*, **30**, 460
- Hinkley, S., Oppenheimer, B. R., Zimmerman, N., et al. 2011, *PASP*, **123**, 74
- Hubeny, I., & Burrows, A. 2007, *ApJ*, **669**, 1248
- Kimble, R. A., MacKenty, J. W., O'Connell, R. W., & Townsend, J. A. 2008, *Proc. SPIE*, **7010**, 70101E-1
- Kirkpatrick, J. D. 2000, in ASP Conf. Ser. 212, From Giant Planets to Cool Stars, ed. C. A. Griffith & M. S. Marley (San Francisco, CA: ASP), 20
- Kirkpatrick, J. D. 2008, in ASP Conf. Ser. 384, 14th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun, ed. G. van Belle (San Francisco, CA: ASP), 85
- Kirkpatrick, J. D., Cushing, M. C., Gelino, C. R., et al. 2011, *ApJS*, **197**, 19
- Kirkpatrick, J. D., Henry, T. J., & McCarthy, D. W., Jr. 1991, *ApJS*, **77**, 417
- Kirkpatrick, J. D., Looper, D. L., Burgasser, A. J., et al. 2010, *ApJS*, **190**, 100
- Kirkpatrick, J. D., Reid, I. N., Liebert, J., et al. 1999, *ApJ*, **519**, 802
- Koekemoer, A. M., Fruchter, A. S., Hook, R. N., & Hack, W. 2002, in The 2002 HST Calibration Workshop: Hubble after the Installation of the ACS and the NICMOS Cooling System, ed. S. Arribas, A. Koekemoer, & B. Whitmore (Baltimore, MD: STScI), 337
- Kumar, S. S. 1963, *ApJ*, **137**, 1121
- Kümmel, M., Walsh, J. R., Pirzkal, N., Kuntschner, H., & Pasquali, A. 2009, *PASP*, **121**, 59
- Lawrence, A., Warren, S. J., Almaini, O., et al. 2007, *MNRAS*, **379**, 1599
- Leggett, S. K., Burningham, B., Saumon, D., et al. 2010, *ApJ*, **710**, 1627
- Leggett, S. K., Cushing, M. C., Saumon, D., et al. 2009, *ApJ*, **695**, 1517
- Leggett, S. K., Marley, M. S., Freedman, R., et al. 2007a, *ApJ*, **667**, 537
- Leggett, S. K., Saumon, D., Marley, M. S., et al. 2007b, *ApJ*, **655**, 1079
- Liu, M. C., et al. 2011, arXiv:1103.0014
- Lodders, K. 1999, *ApJ*, **519**, 793
- Lodders, K., & Fegley, B. 2002, *Icarus*, **155**, 393
- Lucas, P. W., Tinney, C. G., Burningham, B., et al. 2010, *MNRAS*, **408**, L56
- Luhman, K. L., Adame, L., D'Alessio, P., et al. 2005, *ApJ*, **635**, L93
- Luhman, K. L., Burgasser, A. J., & Bochanski, J. J. 2011, *ApJ*, **730**, L9
- Luhman, K. L., et al. 2011, submitted
- Macintosh, B., Graham, J., Palmer, D., et al. 2006, *Proc. SPIE*, **6272**, 62720L-1
- Mainzer, A., Cushing, M. C., Skrutskie, M., et al. 2011, *ApJ*, **726**, 30
- Marley, M. S., & Leggett, S. K. 2009, in The Future of Ultracool Dwarf Science with JWST, ed. H. A. Thronson, M. Stiavelli, & A. Tielens (Dordrecht: Springer), 101
- Marley, M. S., Seager, S., Saumon, D., et al. 2002, *ApJ*, **568**, 335
- Martini, P., Persson, S. E., Murphy, D. C., et al. 2004, *Proc. SPIE*, **5492**, 1653
- McLean, I. S., Becklin, E. E., Bendiksen, O., et al. 1998, *Proc. SPIE*, **3354**, 566
- McLean, I. S., Graham, J. R., Becklin, E. E., et al. 2000, *Proc. SPIE*, **4008**, 1048
- McLean, I. S., McGovern, M. R., Burgasser, A. J., et al. 2003, *ApJ*, **596**, 561
- Morgan, W. W., Keenan, P. C., & Kellman, E. (ed.) 1943, *An Atlas of Stellar Spectra, with an Outline of Spectral Classification* (Chicago, IL: Univ. Chicago Press)
- Rayner, J. T., Cushing, M. C., & Vacca, W. D. 2009, *ApJS*, **185**, 289
- Rayner, J. T., Toomey, D. W., Onaka, P. M., et al. 2003, *PASP*, **115**, 362
- Rodriguez, D. R., Zuckerman, B., Melis, C., & Song, I. 2011, *ApJ*, **732**, L29
- Roellig, T. L., Van Cleve, J. E., Sloan, G. C., et al. 2004, *ApJS*, **154**, 418
- Saumon, D., Geballe, T. R., Leggett, S. K., et al. 2000, *ApJ*, **541**, 374
- Saumon, D., & Marley, M. S. 2008, *ApJ*, **689**, 1327
- Saumon, D., Marley, M. S., Cushing, M. C., et al. 2006, *ApJ*, **647**, 552
- Saumon, D., Marley, M. S., Leggett, S. K., et al. 2007, *ApJ*, **656**, 1136
- Saumon, D., Marley, M. S., Lodders, K., & Freedman, R. S. 2003, in IAU Symp. 211, Brown Dwarfs, ed. E. Martín (Cambridge: Cambridge Univ. Press), 345
- Seifahrt, A., Helling, C., Burgasser, A. J., et al. 2009, in AIP Conf. Ser. 1094, ed. E. Stempels (Melville, NY: AIP), 283
- Simcoe, R. A., Burgasser, A. J., Bernstein, R. A., et al. 2008, *Proc. SPIE*, **7014**, 70140U-1
- Simcoe, R. A., Burgasser, A. J., Bochanski, J. J., et al. 2010, *Proc. SPIE*, **7735**, 773514
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, **131**, 1163
- Skrutskie, M. F., Jones, T., Hinz, P., et al. 2010, *Proc. SPIE*, **7735**, 77353H-1
- Stephens, D. C., Leggett, S. K., Cushing, M. C., et al. 2009, *ApJ*, **702**, 154
- Tokunaga, A. T., Simons, D. A., & Vacca, W. D. 2002, *PASP*, **114**, 180
- van Dam, M. A., Bouchez, A. H., Le Mignant, D., et al. 2006, *PASP*, **118**, 310
- Warren, S. G., & Brandt, R. E. 2008, *J. Geophys. Res. (Atmos.)*, **113**, D14220
- Warren, S. J., Mortlock, D. J., Leggett, S. K., et al. 2007, *MNRAS*, **381**, 1400
- Weights, D. J., Lucas, P. W., Roche, P. F., Pinfield, D. J., & Riddick, F. 2009, *MNRAS*, **392**, 817 (erratum)
- Werner, M. W., Roellig, T. L., Low, F. J., et al. 2004, *ApJS*, **154**, 1
- Whitworth, A., Bate, M. R., Nordlund, Å., Reipurth, B., & Zinnecker, H. 2007, in Protostars and Planets V, ed. B. Reipurth, D. Jewitt, & K. Keil (Tucson, AZ: Univ. Arizona Press), 459
- Whitworth, A. P., & Stamatellos, D. 2006, *A&A*, **458**, 817
- Wilson, J. C., Eikenberry, S. S., Henderson, C. P., et al. 2003, *Proc. SPIE*, **4841**, 451
- Wizinowich, P. L., Le Mignant, D., Bouchez, A. H., et al. 2006, *PASP*, **118**, 297
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, **140**, 1868
- York, D. G., Adelman, J., Anderson, J. E., Jr., et al. 2000, *AJ*, **120**, 1579
- Yurchenko, S. N., Barber, R. J., & Tennyson, J. 2011, *MNRAS*, **413**, 1828